

# Finiteness properties for flat cohomology of varieties over finite fields

Timo Keller

September 16, 2018

## Abstract

We study the  $p$ -primary component of the Tate-Shafarevich group of Abelian schemes over higher dimensional bases over finite fields. We prove that finiteness of the  $p$ -primary component descends by alterations, and prove isogeny invariance of finiteness.

Furthermore, we prove that the first flat cohomology group of a finite flat group scheme and the  $p$ -part of the Brauer group of a product of curves and Abelian varieties over a finite field is finite.

**Keywords:** Abelian varieties of dimension  $> 1$ ; Étale and other Grothendieck topologies and cohomologies; Brauer groups of schemes; Arithmetic ground fields

**MSC 2010:** 11G10, 14F20, 14F22, 14K15

## 1 Introduction

The Tate-Shafarevich group  $\text{III}(\mathcal{A}/X)$  of an Abelian scheme  $\mathcal{A}$  over a base scheme  $X$  is of great importance for the arithmetic of  $\mathcal{A}$ . It classifies everywhere locally trivial  $\mathcal{A}$ -torsors.

In [Kel16], section 4.3, we showed that finiteness of an  $\ell$ -primary component of the Tate-Shafarevich group descends under generically étale  $\ell'$ -alterations. This is used in [Kel18], Corollary 5.11 to prove the finiteness of the Tate-Shafarevich group and an analogue of the Birch-Swinnerton-Dyer conjecture for certain Abelian schemes over higher dimensional bases over finite fields under mild conditions. In [Kel16], section 4.4, we showed that finiteness of an  $\ell$ -primary component of the Tate-Shafarevich group is invariant under étale isogenies.

In this article, we generalise some of our results from [Kel16] for  $\ell^\infty$ -torsion to  $p^\infty$ -torsion: We show:

**Theorem 1** (Corollary 3.5). *The Brauer group of a product of smooth proper curves and Abelian varieties over a finite field is finite.*

**Theorem 2** (Theorem 5.9). *Let  $X$  be a proper integral normal variety over a finite field and  $G/X$  be a finite flat commutative group scheme. Then  $H_{\text{fppf}}^1(X, G)$  is finite.*

This result is used to prove:

**Theorem 3** (Theorem 7.3). *Let  $p$  be a prime. Let  $f : X' \rightarrow X$  be a proper, surjective, generically étale morphism of generical degree prime to  $p$  of integral, normal varieties over a finite field. Let  $X$  be a scheme of characteristic  $p$ . If  $\mathcal{A}$  is an Abelian scheme on  $X$  such that the  $p^\infty$ -torsion of the Tate-Shafarevich group  $\text{III}(\mathcal{A}'/X')$  of  $\mathcal{A}' := f^*\mathcal{A} = \mathcal{A} \times_X X'$  is finite, then the  $p^\infty$ -torsion of the Tate-Shafarevich group  $\text{III}(\mathcal{A}/X)$  is finite.*

**Theorem 4** (Theorem 8.1). *Let  $X/k$  be a proper variety over a finite field  $k$  and  $f : \mathcal{A} \rightarrow \mathcal{A}'$  be an isogeny of Abelian schemes over  $X$ . Let  $p$  be an arbitrary prime. Assume  $f$  étale if  $p \neq \text{char } k$ . Then  $\text{III}(\mathcal{A}/X)[p^\infty]$  is finite if and only if  $\text{III}(\mathcal{A}'/X)[p^\infty]$  is finite.*

**Notation.** For an Abelian group  $A$ , let  $A_{\text{tors}}$  be the torsion subgroup of  $A$ , and  $A_{n\text{-tors}} = A/A_{\text{tors}}$ . Let  $A_{\text{div}}$  be the maximal divisible subgroup of  $A$  and  $A_{n\text{-div}} = A/A_{\text{div}}$ . Denote the cokernel of  $A \xrightarrow{n} A$  by  $A/n$  and its kernel by  $A[n]$ , and the  $p$ -primary subgroup  $\varinjlim_n A[p^n]$  by  $A[p^\infty]$ . Canonical isomorphisms are often denoted by “ $\cong$ ”. We denote Pontryagin duality by  $(-)^D$ , duals of  $R$ -modules or  $\ell$ -adic sheaves by  $(-)^V$ , and duals of Abelian schemes and Cartier duals by  $(-)^t$ . The  $\ell$ -adic valuation  $|\cdot|_\ell$  is taken to be normalised by  $|\ell|_\ell = \ell^{-1}$ . If  $\Gamma$  is a group acting on an Abelian group  $A$ , we denote by  $A^\Gamma$  invariants and by  $A_\Gamma$  coinvariants. By  $X^{(i)}$ , we denote the set of codimension- $i$  points of a scheme  $X$ , and by  $|X|$  the set of closed points. For an Abelian variety  $A$ , we denote its Poincaré bundle by  $\mathcal{P}_A$ . By a variety over a field  $k$  we mean a scheme of finite type over  $k$ .

## 2 Flat cohomology

**Definition 2.1.** An isogeny of group schemes is a surjective group scheme homomorphism with finite kernel.

**Lemma 2.2** (Kummer sequence). Let  $X$  be a scheme of characteristic  $p$  and  $G, G'/X$  be smooth commutative group schemes and assume there is a faithfully flat isogeny  $f : G \rightarrow G'$ . Then there is an exact sequence of sheaves on  $X_{\text{fppf}}$

$$0 \rightarrow \ker(f) \rightarrow G \xrightarrow{f} G' \rightarrow 0$$

This applies in particular to  $G = \mathbf{G}_m$  and  $G = \mathcal{A}$  an Abelian scheme.

*Proof.* Since  $f$  is faithfully flat, in particular surjective, it is an epimorphism of sheaves, compare [Kel18], Lemma 2.10. An isogeny of Abelian schemes is faithfully flat by [Mil86], p. 114f., Proposition 8.1.  $\square$

**Lemma 2.3.** Let  $X$  be a scheme of characteristic  $p$  and  $G/X$  be a smooth commutative group scheme. Then there are comparison isomorphisms

$$H_{\text{fppf}}^i(X, G) = H_{\text{ét}}^i(X, G)$$

*Proof.* See [Mil80], p. 116, Remark III.3.11 (b) and note that the proof given there gives a comparison isomorphism for any topologies between the étale and the flat site.  $\square$

## 3 The $p$ -part of the Brauer group

Let  $X$  be a smooth projective geometrically integral variety over a finite field  $k = \mathbf{F}_q$  of characteristic  $p$  with absolute Galois group  $\Gamma$  topologically generated by the Frobenius  $\text{Frob}$ . By [Kel16], p. 213, Corollary 2.5, one has  $\text{Br}(X) = H_{\text{ét}}^2(X, \mathbf{G}_m) = H_{\text{ét}}^2(X, \mathbf{G}_m)_{\text{tors}}$ , and these equal the corresponding flat cohomology groups by Lemma 2.3.

**Proposition 3.1.** There is a diagram of finitely generated  $\mathbf{Z}_p$ -modules with exact rows and columns

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \text{Pic}(X) \otimes_{\mathbf{Z}} \mathbf{Z}_p & \longrightarrow & (\text{NS}(\bar{X}) \otimes_{\mathbf{Z}} \mathbf{Z}_p)^{\Gamma} & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & H_{\text{fppf}}^1(\bar{X}, \mathbf{Z}_p(1))_{\Gamma} & \longrightarrow & H_{\text{fppf}}^2(X, \mathbf{Z}_p(1)) & \longrightarrow & H_{\text{fppf}}^2(\bar{X}, \mathbf{Z}_p(1))^{\Gamma} \longrightarrow 0 \\
 & & & & \downarrow & & \\
 & & & & T_p \text{Br}(X) & & \\
 & & & & \downarrow & & \\
 & & & & 0 & & 
 \end{array}$$

The group  $H_{\text{fppf}}^1(\bar{X}, \mathbf{Z}_p(1))_{\Gamma}$  is killed by multiplication by  $t(X) := |\text{Pic}(X)_{\text{tors}}| = |\text{Pic}(\bar{X})_{\text{tors}}|^{\Gamma} < \infty$ .

*Proof.* Pass to the projective limit in

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
& & & \text{Pic}(X)/p^n & \longrightarrow & (\text{NS}(\bar{X})/p^n)^\Gamma & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & \text{H}_{\text{fppf}}^1(\bar{X}, \mu_{p^n})_\Gamma & \longrightarrow & \text{H}_{\text{fppf}}^2(X, \mu_{p^n}) & \longrightarrow & \text{H}_{\text{fppf}}^2(\bar{X}, \mu_{p^n})^\Gamma \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & \text{Pic}(\bar{X})[p^n]_\Gamma & & \text{Br}(X)[p^n] & & \\
& & \downarrow & & \downarrow & & \\
& & 0 & & 0 & & 
\end{array}$$

The columns come from the Kummer sequence Lemma 2.2, and the middle row comes from the Hochschild-Serre spectral sequence for the flat topology [Mil80], p. 105, Remark III.2.21 (a). Note that  $\text{H}_{\text{fppf}}^2(X, \mathbf{Z}_p(1))$  is a finitely generated  $\mathbf{Z}_p$ -module by [Ill79], p. 629, Proposition 5.9.

One has  $\text{Pic}(X) = \text{Pic}(\bar{X})^\Gamma$  since the Hochschild-Serre spectral sequence for the flat topology [Mil80], p. 105, Remark III.2.21 (a) gives us an exact sequence

$$0 \rightarrow \text{H}^1(\Gamma, \text{H}_{\text{fppf}}^0(\bar{X}, \mathbf{G}_m)) \rightarrow \text{Pic}(X) \rightarrow \text{Pic}(\bar{X})^\Gamma \rightarrow \text{H}^2(\Gamma, \text{H}_{\text{fppf}}^0(\bar{X}, \mathbf{G}_m)),$$

and  $\text{H}_{\text{fppf}}^0(\bar{X}, \mathbf{G}_m) = \bar{k}^\times$ , and  $\text{H}^1(\Gamma, \bar{k}^\times) = 0$  by Hilbert's theorem 90 and  $\text{H}^2(\Gamma, \bar{k}^\times) = \text{Br}(k) = 0$ . The order of the group  $(\text{Pic}(\bar{X})[p^n])_\Gamma$  equals the order of the group  $(\text{Pic}(\bar{X})[p^n])^\Gamma$  since  $\text{Pic}(\bar{X})[p^n]$  is finite, so its Herbrand quotient equals 1. Hence  $\text{H}_{\text{fppf}}^1(\bar{X}, \mu_{p^n})_\Gamma$  is killed by multiplication with  $t(X)$ , so  $\text{H}_{\text{fppf}}^1(\bar{X}, \mathbf{Z}_p(1))_\Gamma$  is killed by multiplication by  $t(X)$ .  $\square$

**Proposition 3.2.** *There is a diagram of finitely generated  $\mathbf{Q}_p$ -vector spaces with exact rows and columns*

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
& & & \text{Pic}(X) \otimes_{\mathbf{Z}} \mathbf{Q}_p & \xrightarrow{\cong} & (\text{NS}(\bar{X}) \otimes_{\mathbf{Z}} \mathbf{Q}_p)^\Gamma & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & \text{H}_{\text{fppf}}^2(X, \mathbf{Q}_p(1)) & \xrightarrow{\cong} & \text{H}_{\text{fppf}}^2(\bar{X}, \mathbf{Q}_p(1))^\Gamma & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \\
& & V_p \text{Br}(X) & & & & \\
& & \downarrow & & & & \\
& & 0 & & & & 
\end{array}$$

*Proof.* Tensorise the groups in Proposition 3.1 by  $\mathbf{Q}_p$ . Since  $\text{H}_{\text{fppf}}^1(\bar{X}, \mathbf{Z}_p(1))_\Gamma$  is killed by multiplication with  $t(X)$ , it is 0 after tensoring with  $\mathbf{Q}_p$ . The upper horizontal arrow is an isomorphism since  $\text{Pic}(X) = \text{Pic}(\bar{X})^\Gamma$  (see the proof of Proposition 3.1) and  $\text{Pic}(\bar{X}) \otimes_{\mathbf{Z}} \mathbf{Q}_p = \text{NS}(\bar{X}) \otimes_{\mathbf{Z}} \mathbf{Q}_p$  because  $\text{Pic}^0(\bar{X}) = \mathbf{Pic}_{X/k}^0(\bar{k}) = \varinjlim_n \mathbf{Pic}_{X/k}^0(\mathbf{F}_{q^n})$  is torsion as a colimit of finite groups; furthermore,  $(\text{NS}(\bar{X}) \otimes_{\mathbf{Z}} \mathbf{Q}_p)^\Gamma = \text{NS}(\bar{X})^\Gamma \otimes_{\mathbf{Z}} \mathbf{Q}_p$  since  $-\otimes_{\mathbf{Z}} \mathbf{Q}_p$  is exact and  $\text{NS}(\bar{X})$  is a discrete  $\Gamma$ -module and for a finite group  $G$ , one has  $A^G = \varprojlim(A \rightarrow \bigoplus_{g \in G} A)$ .  $\square$

**Corollary 3.3.** *The following are equivalent:*

1. The group  $\text{Br}(X)[p^\infty]$  is finite.

2.  $\mathrm{Br}(X)[p^\infty]_{\mathrm{div}} = 0$  and  $\mathrm{Br}(X)[p^\infty] = \mathrm{Br}(X)[p^\infty]_{\mathrm{n-div}}$ .
3.  $V_p \mathrm{Br}(X) = 0$ .
4.  $\mathrm{NS}(X) \otimes_{\mathbf{Z}} \mathbf{Q}_p = \mathrm{H}_{\mathrm{fppf}}^2(\bar{X}, \mathbf{Q}_p(1))^\Gamma = \mathrm{H}_{\mathrm{fppf}}^2(\bar{X}, \mathbf{Q}_p(1))^{\mathrm{Frob}}$ .
5.  $\dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{fppf}}^2(\bar{X}, \mathbf{Q}_p(1))^\Gamma = \rho(X)$  with  $\rho(X) := \mathrm{rk}_{\mathbf{Z}} \mathrm{NS}(X) = \mathrm{rk}_{\mathbf{Z}} \mathrm{Pic}(X)$ .

One always has  $\dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{fppf}}^2(\bar{X}, \mathbf{Q}_p(1))^\Gamma \geq \rho(X)$ .

**Theorem 3.4.** *Let  $X$  be a product of smooth proper curves and Abelian varieties over a finite field  $k$  of characteristic  $p$ . Then  $\dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{cris}}^2(\bar{X}/W)^\Gamma \otimes \mathbf{Q}_p = \dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{fppf}}^2(X, \mathbf{Q}_p(1)) = \rho(X)$ . Hence  $\mathrm{Br}(X)[p^\infty]$  is finite by Corollary 3.3.*

*Proof.* Note that proper curves and Abelian varieties are projective by [Har83], p. 136, Proposition II.6.7 and [Mil86], p. 113, Theorem 7.1. By Corollary 3.3,

$$\dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{fppf}}^2(X, \mathbf{Q}_p(1)) \geq \rho(X),$$

so it suffices to prove that  $\dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{fppf}}^2(X, \mathbf{Q}_p(1)) \leq \rho(X)$ .

One has for  $\ell \neq p$  prime

$$\rho(X) = \dim_{\mathbf{Q}_\ell} \mathrm{H}_{\mathrm{ét}}^2(\bar{X}, \mathbf{Q}_\ell(1))^\Gamma = \dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{cris}}^2(X/W) \otimes_{\mathbf{Z}} \mathbf{Q}_p,$$

the first equality by the Tate conjecture [Tat66], p. 143, Theorem 4, and the second equality by [KM74], p. 75, Corollary 1.1) since  $X/k$  is smooth projective.

One has  $\mathrm{H}_{\mathrm{ét}}^{2r}(\bar{X}, \mathbf{Q}_\ell(i))^\Gamma = \mathrm{H}_{\mathrm{ét}}^{2r}(X, \mathbf{Q}_\ell(i))$  for  $k$  finite: The Hochschild-Serre spectral sequence

$$\mathrm{H}^p(\Gamma, \mathrm{H}_{\mathrm{ét}}^q(\bar{X}, \mathbf{Q}_\ell(i))) \Rightarrow \mathrm{H}_{\mathrm{ét}}^{p+q}(X, \mathbf{Q}_\ell(i))$$

yields by  $\mathrm{cd}(k) = 1$  a short exact sequence

$$0 \rightarrow \mathrm{H}^1(\Gamma, \mathrm{H}_{\mathrm{ét}}^{q-1}(\bar{X}, \mathbf{Q}_\ell(i))) \rightarrow \mathrm{H}_{\mathrm{ét}}^q(X, \mathbf{Q}_\ell(i)) \rightarrow \mathrm{H}_{\mathrm{ét}}^q(\bar{X}, \mathbf{Q}_\ell(i))^\Gamma \rightarrow 0$$

and  $\mathrm{H}_{\mathrm{ét}}^{q-1}(\bar{X}, \mathbf{Q}_\ell(i))$  is uniquely divisible, so  $\mathrm{H}^1(\Gamma, \mathrm{H}_{\mathrm{ét}}^{q-1}(\bar{X}, \mathbf{Q}_\ell(i))) = 0$ , and the same for  $\mathrm{H}_{\mathrm{fppf}}^q(X, \mathbf{Q}_p(i))$ .

One has  $\dim_{\mathbf{Q}_\ell} \mathrm{H}_{\mathrm{ét}}^2(\bar{X}, \mathbf{Q}_\ell(1)) = \dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{cris}}^2(\bar{X}/W) \otimes_{\mathbf{Z}} \mathbf{Q}_p \geq \dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{fppf}}^2(\bar{X}, \mathbf{Q}_p(1))$ , so

$$\rho(X) = \dim_{\mathbf{Q}_\ell} \mathrm{H}_{\mathrm{ét}}^2(\bar{X}, \mathbf{Q}_\ell(1))^\Gamma \geq \dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{fppf}}^2(X, \mathbf{Q}_p(1)).$$

On the other hand,

$$\dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{fppf}}^2(X, \mathbf{Q}_p(1)) \leq \dim_{\mathbf{Q}_p} \mathrm{H}_{\mathrm{cris}}^2(X/W) \otimes_{\mathbf{Z}} \mathbf{Q}_p$$

by [Ill79], p. 627, Théorème 5.5 (5.5.3) or p. 631, Théorème 5.14.

Tate's conjecture [Tat66] implies that the Frobenius action on the cohomology is semisimple, hence  $\dim_{\mathbf{Q}_\ell} \mathrm{H}_{\mathrm{ét}}^2(\bar{X}, \mathbf{Q}_\ell(1))^\Gamma$  is equal to the multiplicity of  $1 - X$  as a factor of the characteristic polynomial of the Frobenius, which is independent of  $\ell$ . This also works for crystalline cohomology (which is a Weil cohomology theory).  $\square$

**Corollary 3.5.** *The Brauer group of a product of smooth proper curves and Abelian varieties over a finite field is finite.*

*Proof.* Combine Theorem 3.4 with [Zar83], p. 214, Corollary 2.3.5 and use that the Brauer group of a regular scheme is torsion.  $\square$

## 4 The $p$ -part of the Brauer group of an Abelian variety over a finite field

In this section, we prove a formula for the order of  $\mathrm{Br}(A)[p^\infty]$  for an Abelian variety  $A$  over a finite field of characteristic  $p$ .

**Lemma 4.1.** *Let  $k$  be a perfect field,  $W = W(k)$  the ring of Witt vectors and  $A/k$  be an Abelian variety. Then  $H_{\mathrm{cris}}^i(A/W) = \bigwedge^i H_{\mathrm{cris}}^1(A/W)$ .*

*Proof.* See [Ill79], p. 651, (7.1.1). □

**Lemma 4.2.** *Let  $k$  be a finite field of characteristic  $p$  and  $A/k$  be an Abelian variety. Then the action of  $\Gamma = \mathrm{Gal}(\bar{k}/k)$  on  $T_p A$  is semisimple.*

*Proof.* Let  $\ell \neq p$  be prime. The minimal polynomial of the Frobenius acting on  $T_\ell A$  is defined over  $\mathbf{Q}$  since it equals the radical of the characteristic polynomial, which is defined over  $\mathbf{Q}$  by [Mil86], p. 125, Proposition 12.9.

By [Tat66], the Frobenius acts semisimply on  $T_\ell A$  for  $\ell \neq p$  prime, and hence its minimal polynomial is square-free. Since  $\mathrm{End}_k(A) \otimes_{\mathbf{Z}} \mathbf{Q}_\ell \rightarrow \mathrm{End}_{\mathbf{Q}_\ell}(T_\ell A)$  is injective, it also satisfies this square-free polynomial in  $\mathrm{End}_k(A) \otimes_{\mathbf{Z}} \mathbf{Q} \subset \mathrm{End}_k(A) \otimes_{\mathbf{Z}} \mathbf{Q}_\ell$ , so it is semisimple. Hence it also acts semisimply on  $T_p A$  as there is an injection  $\mathrm{End}_k(A) \otimes_{\mathbf{Z}} \mathbf{Q}_p \hookrightarrow \mathrm{End}_{\mathbf{Q}_p}(T_p A)$ . □

## 5 Finiteness theorems for flat and syntomic cohomology over finite fields

The aim of this section is to show that  $H_{\mathrm{fppf}}^1(X, G)$  is finite for  $X$  a normal proper variety over a finite field and  $G/X$  a finite flat group scheme.

The proof is by reduction to the case of a finite flat simple group scheme over an algebraically closed field, which is isomorphic to  $\mathbf{Z}/p$ ,  $\mu_p$  or  $\alpha_p$ .

We use the interpretation of  $H_{\mathrm{fppf}}^1(X, G)$  as  $G$ -torsors on  $X$  [Mil80], p. 124, Proposition III.4.7 since  $G/X$  is affine. We also use de Jong's alteration theorem [de 96], p. 66, Theorem 4.1.

**Lemma 5.1.** *Let  $X$  be a Noetherian integral scheme with function field  $K(X)$  and  $U \subseteq X$  dense open. Then there is an exact sequence*

$$1 \rightarrow \mathbf{G}_m(X) \rightarrow \mathbf{G}_m(U) \rightarrow \bigoplus_{D \in (X \setminus U)^{(1)}} \mathbf{Z}[D] \rightarrow \mathrm{Cl}(X) \rightarrow \mathrm{Cl}(U) \rightarrow 0.$$

*Proof.* The assumptions imply that there is a commutative diagram with exact rows

$$\begin{array}{ccccccccc} 1 & \longrightarrow & \mathcal{O}_X(X)^\times & \longrightarrow & K(X)^\times & \longrightarrow & \mathrm{Div}(X) & \longrightarrow & \mathrm{Cl}(X) & \longrightarrow & 0 \\ & & \downarrow & & \parallel & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \mathcal{O}_X(U)^\times & \longrightarrow & K(U)^\times & \longrightarrow & \mathrm{Div}(U) & \longrightarrow & \mathrm{Cl}(U) & \longrightarrow & 0. \end{array}$$

A diagram chase yields the result. □

**Corollary 5.2.** *Let  $X$  be a Noetherian integral regular scheme and let  $U \subseteq X$  be dense open. Then there is an exact sequence*

$$1 \rightarrow \mathbf{G}_m(X) \rightarrow \mathbf{G}_m(U) \rightarrow \bigoplus_{D \in (X \setminus U)^{(1)}} \mathbf{Z}[D] \rightarrow \mathrm{Pic}(X) \rightarrow \mathrm{Pic}(U) \rightarrow 0.$$

*Proof.* By the assumptions,  $\mathrm{Cl}(X) = \mathrm{Pic}(X)$  and  $\mathrm{Cl}(U) = \mathrm{Pic}(U)$ . □

**Corollary 5.3.** *Let  $X/\mathbf{F}_q$  be an integral Noetherian regular proper variety and let  $j : U \hookrightarrow X$  be the inclusion of an open subscheme of  $X$ . Then  $H_{\mathrm{fppf}}^1(U, \mu_{p^n})$  is finite for all  $n$  and any prime  $p$ .*

*Proof.* The Kummer sequence Lemma 2.2 on  $U_{\text{fppf}}$  together with  $\text{Pic}(U) = H_{\text{fppf}}^1(U, \mathbf{G}_{m,U})$  by Lemma 2.3 yields the exact sequence

$$1 \rightarrow \mathbf{G}_m(U)/p^n \rightarrow H_{\text{fppf}}^1(U, \mu_{p^n}) \rightarrow \text{Pic}(U)[p^n] \rightarrow 0.$$

Since  $\mathbf{G}_m(X) = \Gamma(X, \mathbf{G}_m)^\times$  is finite by the coherence theorem since  $X/\mathbf{F}_q$  is proper and  $\mathbf{F}_q$  is finite, and since  $\text{Pic}(X)$  is finitely generated since its sits in a short exact sequence  $0 \rightarrow \text{Pic}^0(X) \rightarrow \text{Pic}(X) \rightarrow \text{NS}(X) \rightarrow 0$  and  $\text{Pic}^0(X)$  is finite since it is the group of rational points of an Abelian variety over a finite field and  $\text{NS}(X)$  is always finitely generated by [Mil80], p. 215, Theorem V.3.25, by Corollary 5.2 and the finiteness of  $(X \setminus U)^{(1)}$ , this exact sequence gives the finiteness of  $\mathbf{G}_m(U)/p^n$  and of  $\text{Pic}(U)[p^n]$ .  $\square$

**Lemma 5.4.** *Let  $X$  be a normal integral scheme and  $G/X$  be a finite flat group scheme. If  $T$  is a  $G$ -torsor on  $X$  trivial over the generic point of  $X$ , then  $T$  is trivial. Hence,  $H_{\text{fppf}}^1(X, G) \rightarrow H_{\text{fppf}}^1(K(X), G)$  is injective, and if  $f : Y \rightarrow X$  is birational,  $f^* : H_{\text{fppf}}^1(X, G) \rightarrow H_{\text{fppf}}^1(Y, G)$  is injective.*

*Proof.* Since  $T$  is trivial over the generic point of  $X$ , generically, there is a section of  $\pi : T \rightarrow X$ . This extends to a rational map  $\sigma : X \dashrightarrow T$ . Take the schematic closure  $i : X' \hookrightarrow T$  of  $\sigma$ . The composition  $\pi \circ i : X' \rightarrow T \rightarrow X$  is birational and finite (as a composition of a closed immersion and a finite morphism). By [GW10], p. 358, Corollary 12.88, since  $X$  is normal,  $X' \rightarrow X$  is an isomorphism. Hence  $\sigma$  is a section of  $\pi$ , so  $T/X$  is trivial.  $\square$

**Lemma 5.5.** *Let  $X$  be a proper variety over a finite field and  $Y/X$  be a finite flat scheme. Let  $Z/X$  be proper. Then  $Y(Z)$  is finite.*

*Proof.* Since  $\text{Mor}_X(Z, Y) = \text{Mor}_Z(Z, Y \times_X Z)$ , one can assume  $Z = X$ . So we have to show that there are only finitely many sections to  $\pi : Y \rightarrow X$ . Such a section corresponds to an  $\mathcal{O}_X$ -algebra map  $\pi_* \mathcal{O}_Y \rightarrow \mathcal{O}_X$ . But  $H_{\text{Zar}}^0(X, \mathcal{H}\text{om}_X(\pi_* \mathcal{O}_Y, \mathcal{O}_X))$  is finite by the coherence theorem as it is a finite dimensional vector space over a finite field.  $\square$

**Lemma 5.6.** *Let  $Y \rightarrow X$  be an alteration of proper integral varieties with  $X$  normal, and  $G/X$  be a finite flat commutative group scheme. Then  $\ker(H_{\text{fppf}}^1(X, G) \rightarrow H_{\text{fppf}}^1(Y, G))$  is finite. Hence  $H_{\text{fppf}}^1(X, G)$  is finite if  $H_{\text{fppf}}^1(Y, G)$  is.*

*Proof.* If  $Y \rightarrow X$  is a blow-up, the kernel is trivial by Lemma 5.4 since a blow-up is birational. Hence the statement holds for blow-ups.

Since a normalisation morphism of integral schemes is birational [Liu06], p. 120, Proposition 4.1.22, one can assume  $X'$  normal.

By [RG71], p. 37, Théorème 5.2.2, there is a blow-up  $X' \rightarrow X$  such that  $Y' := Y \times_X X'$  is flat over  $X'$ . There is a commutative diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & \ker(H_{\text{fppf}}^1(X, G) \rightarrow H_{\text{fppf}}^1(Y, G)) & \longrightarrow & H_{\text{fppf}}^1(X, G) \\ & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \ker(H_{\text{fppf}}^1(X', G) \rightarrow H_{\text{fppf}}^1(Y', G)) & \longrightarrow & H_{\text{fppf}}^1(X', G) \end{array}$$

By the snake lemma, since  $\ker(H_{\text{fppf}}^1(X, G) \rightarrow H_{\text{fppf}}^1(X', G))$  is finite as  $X' \rightarrow X$  is a blow-up,  $\ker(H_{\text{fppf}}^1(X, G) \rightarrow H_{\text{fppf}}^1(Y, G))$  is finite if we can show that  $\ker(H_{\text{fppf}}^1(X', G) \rightarrow H_{\text{fppf}}^1(Y', G))$  is finite. Hence, we can assume  $Y \rightarrow X$  finite flat.

Let  $T \rightarrow X$  be in the kernel, i.e. it is a torsor trivial on  $Y$ . Choose a section  $\sigma : Y \rightarrow T \times_X Y$ . Since  $T \times_X (Y \times_X Y) \rightarrow Y \times_X Y$  is a  $G$ -torsor, one can take

$$\tau := \sigma \circ \text{pr}_0 - \sigma \circ \text{pr}_1 \in G(Y \times_X Y).$$

The section  $\tau$  corresponds to the isomorphism class of the  $G$ -torsor  $T$  by descent theory for the fppf covering  $\{Y \rightarrow X\}$ , but by Lemma 5.5,  $G(Y \times_X Y)$  is finite.  $\square$

**Lemma 5.7.** *Let  $X$  be an integral scheme with function field  $K$  and  $G/X$  be a finite flat group scheme. Let  $H_K \hookrightarrow G_K$  be a finite flat group scheme. Then there is a blow-up  $\tilde{X}/X$  such that  $H_K$  extends to a finite flat subgroup scheme of  $G \times_X \tilde{X}$ .*

*Proof.* Let  $H \hookrightarrow G$  be the schematic closure of  $H_K \hookrightarrow G$ . The morphism  $H \rightarrow G \rightarrow X$  is finite as a composition of a closed immersion and a finite morphism. By [RG71], p. 37, Théorème 5.2.2, there is a blow-up  $X' \rightarrow X$  such that  $H' := H \times_X X' \rightarrow X'$  is flat. Then,  $H'$  is the schematic closure of  $H_K \hookrightarrow G' := G \times_X X'$ . So one can assume  $H/X$  finite flat.

Let  $Y \rightarrow X$  be finite flat. Since the morphism is affine, locally, one has the diagram

$$\begin{array}{ccc} A^C & \longrightarrow & A \otimes_R \text{Quot}(R) \\ \uparrow & & \uparrow \\ R^C & \longrightarrow & \text{Quot}(R) \end{array}$$

Here, the upper horizontal arrow is injective by flatness of  $R \rightarrow A$ . Hence  $Y$  is the schematic closure of  $Y_K$  in  $Y$ .

By flatness, the schematic closure of  $H_K \times_K H_K$  in  $G \times_X G$  is  $H \times_X H$ . By the universal property of the schematic closure [GW10], p. 251, (10.8), one has the factorisation

$$\begin{array}{ccc} H_K \times_K H_K & \xrightarrow{\mu} & H_K \\ \downarrow & & \downarrow \\ H \times_X H & \xrightarrow{\mu} & H \\ \downarrow & & \downarrow \\ G \times_X G & \xrightarrow{\mu} & G, \end{array}$$

for the multiplication  $\mu$ , and similar for the inverse and unit section.  $\square$

**Lemma 5.8.** *Let  $X$  be a proper integral variety over a field and  $G/X$  be a finite flat commutative group scheme. After an alteration  $X' \rightarrow X$ , there exists a filtration of  $G$  by finite flat group schemes with subquotients of prime order.*

*Proof.* Over the algebraic closure of the function field of  $X$ , there is such a filtration since the only simple objects in the category of finite flat group schemes of  $p$ -power order are  $\mu_p$ ,  $\mathbf{Z}/p$  and  $\alpha_p$ . Since everything is of finite presentation, these are defined over a finite extension of the function field [GW10], p. 269, Corollary 10.79. Now take the normalisation in this finite extension of function fields and use Lemma 5.7.  $\square$

**Theorem 5.9.** *Let  $X$  be a proper integral normal variety over a finite field and  $G/X$  be a finite flat commutative group scheme. Then  $H_{\text{fppf}}^1(X, G)$  is finite.*

*Proof.* By Lemma 5.8, Lemma 5.6 and the long exact cohomology sequence one can assume  $G$  of prime order  $p$  (since the case of  $G/X$  étale is easily dealt with). Since then  $G$  is simple by [Sha86], p. 38, and since  $F \circ V = [p] = 0$  by [Sha86], p. 62 and [Mum70], p. 141, either  $V = 0$  or  $F = 0$  on  $G$ .

If  $V = 0$ , by [de 93], p. 93, Proposition 2.2, there is a short exact sequence

$$0 \rightarrow G \rightarrow \mathcal{L} \rightarrow \mathcal{M} \rightarrow 0$$

with vector bundles  $\mathcal{L}, \mathcal{M}$ . By the coherence theorem, as  $X$  is proper and lives over a finite ground field, and by comparison of Zariski and fppf cohomology [Mil80], p. 114, Proposition III.3.7, the long exact cohomology sequence shows that  $H_{\text{fppf}}^2(X, G)$  is finite.

If  $F = 0$ , after replacing  $X$  by an alteration by Lemma 5.6 as in the proof of Lemma 5.8, one can assume that  $G$  is isomorphic to  $\mu_p$  over the generic point. Since for  $Y, Z/X$  of finite presentation such that  $Y_K \cong Z_K$ , there is a non-empty open subscheme  $U \hookrightarrow X$  such that  $Y_U \cong Z_U$ , there is a non-empty open subscheme  $U \hookrightarrow X$  such that  $G_U \cong \mu_{p,U}$ . There is an alteration  $f : X' \rightarrow X$  such that  $X'$  is regular. By Corollary 5.3,  $H_{\text{fppf}}^1(f^{-1}(U), \mu_p)$  is finite. By Lemma 5.4,  $H_{\text{fppf}}^1(X', G \times_X X')$  is finite, so by Lemma 5.6,  $H_{\text{fppf}}^1(X, G)$  is finite.  $\square$

## 6 Vanishing of étale cohomology with supports of Abelian schemes

This is a complement to the vanishing condition [Kel16], p. 226, (4.4), which is proven there only for Jacobians of curves, see [Kel16], p. 229, Lemma 4.10.

**Theorem 6.1.** *Let  $X$  be a regular integral Noetherian separated scheme and  $G/X$  be a finite étale commutative group scheme of order invertible on  $X$ . Let  $Z \hookrightarrow X$  be a closed subscheme of codimension  $\geq 2$ . Then  $H_Z^i(X, G) = 0$  for  $i \leq 2$  (étale cohomology with supports in  $Z$ ).*

*Proof.* Let  $U = X \setminus Z$ . One has a long exact cohomology sequence

$$\dots \rightarrow H^{i-1}(X, G) \rightarrow H^{i-1}(U, G) \rightarrow H_Z^i(X, G) \rightarrow H^i(X, G) \rightarrow H^i(U, G) \rightarrow \dots,$$

so one has to prove that  $H^i(X, G) \rightarrow H^i(U, G)$  is an isomorphism for  $i = 0, 1$  and injective for  $i = 2$ .

For  $i = 0$ , the claim  $H_Z^0(X, G) = 0$  is equivalent to the injectivity of

$$H^0(X, G) \rightarrow H^0(X \setminus Z, G),$$

which is clear from [Har83], p. 105, Exercise II.4.2 since  $G/X$  is separated,  $X$  is reduced and  $X \setminus Z \hookrightarrow X$  is dense.

For  $i = 1$  the claim  $H_Z^1(X, G) = 0$  is equivalent to

$$H^0(X, G) \rightarrow H^0(X \setminus Z, G)$$

being surjective and

$$H^1(X, G) \rightarrow H^1(X \setminus Z, G)$$

being injective. The surjectivity of  $H^0(X, G) \rightarrow H^0(X \setminus Z, G)$  follows e. g. from

**Theorem 6.2.** *Let  $S$  be a normal Noetherian base scheme, and let  $u : Z \dashrightarrow G$  be an  $S$ -rational map from a smooth  $S$ -scheme  $Z$  to a smooth and separated  $S$ -group scheme  $G$ . Then, if  $u$  is defined in codimension  $\leq 1$ , it is defined everywhere.*

*Proof.* See [BLR90], p. 109, Theorem 1. □

For the injectivity of  $H^1(X, G) \rightarrow H^1(X \setminus Z, G)$ : If a principal homogeneous space  $P/X$  for  $G/X$  is trivial over  $X \setminus Z$ , then it is trivial over  $X$ : The trivialisation over  $X \setminus Z$  gives a rational map from  $X$  to the principal homogeneous space and any such map (with  $X$  a regular scheme) extends to a morphism by Theorem 6.2.

For the surjectivity of  $H^1(X, G) \rightarrow H^1(X \setminus Z, G)$ : This means that any principal homogeneous space  $P/(X \setminus Z)$  extends to a principal homogeneous space  $\bar{P}/X$ . By [Mil80], p. 123, Corollary III.4.7, we have  $\mathrm{PHS}(G/X) \xrightarrow{\sim} H^1(X_{\mathrm{fl}}, G)$  (Čech cohomology) since  $G/X$  is affine. Since  $G/X$  is smooth, [Mil80], p. 123, Remark III.4.8 (a) shows that we can take étale cohomology as well, and by [Mil80], p. 101, Corollary III.2.10, one can take derived functor cohomology instead of Čech cohomology. By Zariski-Nagata purity [SGA1], Exp. X, Corollaire 3.3, one can extend this to a  $\bar{P}/X$ , for which we have to show that it represents an element of  $H^1(X, G)$ , i. e. that it is a  $G$ -torsor.

Now we need to show that if  $P/(X \setminus Z)$  is an  $G|_{X \setminus Z}$ -torsor and  $\bar{P}$  an extension of  $P$  to a finite étale covering of  $X$ , then  $\bar{P}/X$  is also an  $G$ -torsor. For this, we use the following

**Theorem 6.3.** *Let  $S$  be a connected scheme,  $G \rightarrow S$  a finite flat group scheme, and  $X \rightarrow S$  a scheme over  $S$  equipped with a left action  $\rho : G \times_S X \rightarrow X$ . These data define a  $G$ -torsor over  $S$  if and only if there exists a finite locally free surjective morphism  $Y \rightarrow S$  such that  $X \times_S Y \rightarrow Y$  is isomorphic, as a  $Y$ -scheme with  $G \times_S Y$ -action, to  $G \times_S Y$  acting on itself by left translations.*

*Proof.* See [Sza09], p. 171, Lemma 5.3.13. □

That  $P/(X \setminus Z)$  is an  $G|_{X \setminus Z}$ -torsor amounts to saying that there is an operation

$$G|_{X \setminus Z} \times_{X \setminus Z} P \rightarrow P$$

as in the previous Theorem 6.3. Since this is étale locally isomorphic to the canonical action

$$G|_{X \setminus Z} \times_{X \setminus Z} G|_{X \setminus Z} \xrightarrow{\mu} G|_{X \setminus Z}$$

which is finite étale, by faithfully flat descent the operation defines an étale covering, so extends by Zariski-Nagata purity uniquely to an étale covering  $H \rightarrow X$ , which by uniqueness has to be isomorphic to  $G \times_X \bar{P} \rightarrow \bar{P}$ . Now one has to check the condition in Theorem 6.3.

There is a finite étale Galois covering  $X'/X$  with Galois group  $G$  such that  $G \times_X X'$  is isomorphic to a direct sum of  $\mu_n$  with  $n$  invertible on  $X$ . The Leray spectral sequence with supports  $H^p(G, H_Z^q(X', G \times_X X')) \Rightarrow H_Z^{p+q}(X, G)$  from [Kel16], p. 228, Theorem 4.9, so it suffices to show  $H_Z^q(X', G \times_X X') = 0$  for  $q = 0, 1, 2$ . Hence one can assume  $G \cong \mu_n$  for  $n$  invertible on  $X$ .

One has an injection  $\text{Br}(X) \hookrightarrow \text{Br}(K(X))$  with  $K(X)$  the function field of  $X$  and  $\text{Br}(X) \rightarrow \text{Br}(U) \rightarrow \text{Br}(K(X))$ , so  $\text{Br}(X) \rightarrow \text{Br}(U)$  is injective. By the hypotheses on  $X$  and since the codimension of  $Z$  in  $X$  is  $\geq 2$ , by Corollary 5.2, there is a restriction isomorphism  $\text{Pic}(X) \xrightarrow{\sim} \text{Pic}(U)$ . Hence the snake lemma applied to the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Pic}(X)/n & \longrightarrow & H^2(X, \mu_n) & \longrightarrow & \text{Br}(X)[n] \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \text{Pic}(U)/n & \longrightarrow & H^2(U, \mu_n) & \longrightarrow & \text{Br}(U)[n] \longrightarrow 0 \end{array}$$

gives that  $H^2(X, \mu_n) \rightarrow H^2(U, \mu_n)$  is injective, so  $H_Z^2(X, \mu_n) = 0$ .  $\square$

**Corollary 6.4.** *Let  $X$  be a regular integral Noetherian separated scheme and  $\mathcal{A}/X$  be an Abelian variety. Let  $Z \hookrightarrow X$  be a closed subscheme of codimension  $\geq 2$ . Then  $H_Z^i(X, \mathcal{A})$  is torsion for all  $i$ ,  $= 0$  for  $i = 0$  and  $H_Z^i(X, \mathcal{A})[p^\infty] = 0$  for  $i = 0, 1, 2$  and  $p$  invertible on  $X$ .*

*Proof.* By [Kel16], p. 224, Proposition 4.1,  $H^i(X, \mathcal{A})$  is torsion for  $i > 0$ . The Kummer exact sequence  $0 \rightarrow \mathcal{A}[n] \rightarrow \mathcal{A} \rightarrow \mathcal{A} \rightarrow 0$  for  $n$  invertible on  $X$  yields a surjection

$$H_Z^i(X, \mathcal{A}[n]) \twoheadrightarrow H_Z^i(X, \mathcal{A})[n],$$

so it suffices to show that  $H_Z^i(X, \mathcal{A}[n]) = 0$  for  $i = 1, 2$ . But this is Theorem 6.1. The triviality  $H_Z^0(X, \mathcal{A}) = 0$  is equivalent to the injectivity of

$$H^0(X, \mathcal{A}) \rightarrow H^0(X \setminus Z, \mathcal{A}),$$

which is clear from [Har83], p. 105, Exercise II.4.2 since  $\mathcal{A}/X$  is separated,  $X$  is reduced and  $X \setminus Z \hookrightarrow X$  is dense.  $\square$

## 7 Descent of finiteness of III, the $p$ -part

In this section, we extend [Kel16], p. 238, Theorem 4.29 to  $p^\infty$ -torsion.

**Lemma 7.1.** *Let  $\mathcal{A}/X$  be an Abelian scheme over a proper variety  $X$  over a finite field of characteristic  $p$ . Then  $\text{III}(\mathcal{A}/X)[p^\infty]$  is cofinitely generated.*

Recall that  $\text{III}(\mathcal{A}/X)$  was defined as  $H_{\text{ét}}^1(X, \mathcal{A})$  in [Kel16], p. 225, Definition 4.2.

*Proof.* The long exact cohomology sequence associated to the Kummer sequence Lemma 2.2 gives us a surjection

$$H_{\text{fppf}}^1(X, \mathcal{A}[p^n]) \twoheadrightarrow H_{\text{fppf}}^1(X, \mathcal{A})[p^n] \rightarrow 0$$

Now, since  $\mathcal{A}/X$  is a smooth group scheme, Lemma 2.3 gives us an isomorphism  $H_{\text{fppf}}^1(X, \mathcal{A}) = H_{\text{ét}}^1(X, \mathcal{A})$ , which by definition equals  $\text{III}(\mathcal{A}/X)$ . By Theorem 5.9,  $H_{\text{fppf}}^1(X, \mathcal{A}[p^n])$  is finite since  $X/\mathbf{F}_q$  is proper. From this, one sees that  $H_{\text{ét}}^1(X, \mathcal{A})[p]$  is finite. Hence  $\text{III}(\mathcal{A}/X)[p^\infty]$  is cofinitely generated by [Kel18], Lemma 2.38.  $\square$

**Lemma 7.2.** *Let  $f : X' \rightarrow X$  be a finite étale morphism of constant degree  $d$  and let  $\mathcal{F}$  be an fppf sheaf on  $X$ . Then there is a trace map  $\text{Tr}_f : f_* f^* \mathcal{F} \rightarrow \mathcal{F}$ , functorial in  $\mathcal{F}$ , such that  $\varphi \mapsto \text{Tr}_f \circ f_*(\varphi)$  is an isomorphism  $\text{Hom}_{X'}(\mathcal{F}', f^* \mathcal{F}) \rightarrow \text{Hom}_X(\pi_* \mathcal{F}', \mathcal{F})$  for any fppf sheaf  $\mathcal{F}'$  on  $X'$ . Thus,  $f_* = f!$ , that is,  $f_*$  is left adjoint to  $f^*$ , and  $\text{Tr}_f$  is the adjunction map. The composites*

$$\mathcal{F} \rightarrow f_* f^* \mathcal{F} \xrightarrow{\text{Tr}_f} \mathcal{F} \quad \text{and} \quad H_{\text{fppf}}^r(X, \mathcal{F}) \xrightarrow{f^*} H_{\text{fppf}}^r(X', f^* \mathcal{F}) \xrightarrow{\text{can}} H_{\text{fppf}}^r(X, f_* f^* \mathcal{F}) \xrightarrow{\text{Tr}_f} H_{\text{fppf}}^r(X, \mathcal{F})$$

are multiplication by  $d$ .

*Proof.* As in [Mil80], p. 168, Lemma V.1.12.  $\square$

**Theorem 7.3.** *Let  $p$  be a prime. Let  $f : X' \rightarrow X$  be a proper, surjective, generically étale morphism of generical degree prime to  $p$  of integral, normal varieties over a finite field. Let  $X$  be a scheme of characteristic  $p$ . If  $\mathcal{A}$  is an Abelian scheme on  $X$  such that the  $p^\infty$ -torsion of the Tate-Shafarevich group  $\text{III}(\mathcal{A}'/X')$  of  $\mathcal{A}' := f^*\mathcal{A} = \mathcal{A} \times_X X'$  is finite, then the  $p^\infty$ -torsion of the Tate-Shafarevich group  $\text{III}(\mathcal{A}/X)$  is finite.*

*Proof.* The same proof as in [Kel16], p. 238, Theorem 4.29 works, one only needs  $\text{III}(\mathcal{A}/X)[p^\infty]$  to be cofinitely generated in Step 2, which is Lemma 7.1. The trace morphism in Step 3 for fppf cohomology comes from Lemma 7.2. Note that the proof given there does not need the regularity of  $X, X'$  and that varieties over a field are excellent by [Liu06], p. 343, Corollary 2.40 (a).  $\square$

## 8 Isogeny invariance of finiteness of $\text{III}$ , the $p$ -part

In this section, we extend [Kel16], p. 240, Theorem 4.31 to  $p^\infty$ -torsion.

**Theorem 8.1.** *Let  $X/k$  be a proper variety over a finite field  $k$  and  $f : \mathcal{A} \rightarrow \mathcal{A}'$  be an isogeny of Abelian schemes over  $X$ . Let  $p$  be an arbitrary prime. Assume  $f$  étale if  $p \neq \text{char } k$ . Then  $\text{III}(\mathcal{A}/X)[p^\infty]$  is finite if and only if  $\text{III}(\mathcal{A}'/X)[p^\infty]$  is finite.*

*Proof.* In the case where  $\ell$  is invertible on  $X$  and  $f$  is étale (i. e., of degree invertible on  $X$ ), this is [Kel16], p. 240, Theorem 4.31.

Now assume  $p = \text{char } k$ . The short exact sequence of flat sheaves Lemma 2.2 yields an exact sequence in cohomology

$$\mathrm{H}_{\text{fppf}}^1(X, \ker(f)) \rightarrow \mathrm{H}_{\text{fppf}}^1(X, \mathcal{A}) \xrightarrow{f} \mathrm{H}_{\text{fppf}}^1(X, \mathcal{A}')$$

and note that  $\mathrm{H}_{\text{fppf}}^1(X, \mathcal{A}) = \mathrm{H}_{\text{ét}}^1(X, \mathcal{A}) = \text{III}(\mathcal{A}/X)$  by Lemma 2.3 since  $\mathcal{A}/X$  is smooth, and that  $\mathrm{H}_{\text{fppf}}^1(X, \ker(f))$  is finite by Theorem 5.9. Note that all groups are torsion (the Tate-Shafarevich groups by [Kel16], p. 224, Proposition 4.1), hence the sequence stays exact after taking  $p^\infty$ -torsion. So  $\text{III}(\mathcal{A}/X)[p^\infty]$  is finite if  $\text{III}(\mathcal{A}'/X)[p^\infty]$  is.

For the converse, note that by [Kel18], Proposition 2.19 there is a polarisation  $\lambda : \mathcal{A}^t \rightarrow \mathcal{A}$ . Hence, the argument above for  $\lambda$  and  $\lambda^t$  implies that  $\text{III}(\mathcal{A}^t/X)[p^\infty]$  is finite iff  $\text{III}(\mathcal{A}/X)[p^\infty]$  is, and analogously for  $\text{III}(\mathcal{A}'/X)[p^\infty]$ . Taking the dual Kummer sequence  $0 \rightarrow \ker(f^t) \rightarrow \mathcal{A}'^t \rightarrow \mathcal{A}^t \rightarrow 0$  yields an exact sequence

$$\mathrm{H}_{\text{fppf}}^1(X, \ker(f^t)) \rightarrow \text{III}(\mathcal{A}'^t/X) \rightarrow \text{III}(\mathcal{A}^t/X).$$

By the same argument as above,  $\text{III}(\mathcal{A}'^t/X)[p^\infty]$  is finite if  $\text{III}(\mathcal{A}^t/X)[p^\infty]$  is if  $\text{III}(\mathcal{A}/X)[p^\infty]$  is. So  $\text{III}(\mathcal{A}'/X)[p^\infty]$  is finite.  $\square$

**Acknowledgements.** I thank Jean-Louis Colliot-Thélène, Aise Johan de Jong, Ulrich Görtz and Uwe Jannsen, and Ariyan Javanpeykar, darx and David Loeffler from mathoverflow.

## References

- [BLR90] BOSCH, Siegfried; LÜTKEBOHMERT, Werner and RAYNAUD, Michel: *Néron models*. Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, **21**. Berlin etc.: Springer-Verlag, x, 325 p. 1990.
- [de 93] DE JONG, Aise J.: *Finite locally free group schemes in characteristic  $p$  and Dieudonné modules*. In: Invent. Math., **114**(1) (1993), 89–137.
- [de 96] ———: *Smoothness, semi-stability and alterations*. In: Publ. Math., Inst. Hautes Étud. Sci., **83** (1996), 51–93.
- [SGA1] GROTHENDIECK, Alexandre: *Séminaire de Géométrie Algébrique du Bois Marie – 1960–61 – Revêtements étales et groupe fondamental – (SGA 1)*. (Lecture notes in mathematics **224**), Berlin; New York: Springer-Verlag, xxii+447 1971.
- [GW10] GÖRTZ, Ulrich and WEDHORN, Torsten: *Algebraic geometry I. Schemes. With examples and exercises*. Advanced Lectures in Mathematics. Wiesbaden: Vieweg+Teubner, vii, 615 p. 2010.
- [Har83] HARTSHORNE, Robin: *Algebraic geometry. Corr. 3rd printing*. Graduate Texts in Mathematics, **52**. New York-Heidelberg-Berlin: Springer-Verlag, XVI, 496 p. 1983.
- [Ill79] ILLUSIE, Luc: *Complexe de De Rham-Witt et cohomologie cristalline*. In: Ann. Sci. Éc. Norm. Supér. (4), **12** (1979), 501–661.

- [Kel16] KELLER, Timo: *On the Tate-Shafarevich group of Abelian schemes over higher dimensional bases over finite fields*. In: *manuscripta math.*, **150** (2016), 211–245. URL <https://www.timokeller.name/TateShafarevich.pdf>.
- [Kel18] ———: *On an analogue of the conjecture of Birch and Swinnerton-Dyer for Abelian schemes over higher dimensional bases over finite fields*. Preprint, 2018. URL <https://www.timokeller.name/BSDI.pdf>.
- [KM74] KATZ, Nicholas M. and MESSING, William: *Some consequences of the Riemann hypothesis for varieties over finite fields*. In: *Invent. Math.*, **23** (1974), 73–77.
- [Liu06] LIU, Qing: *Algebraic geometry and arithmetic curves*. *Transl. by Reinie Ern e*. Oxford Graduate Texts in Mathematics **6**. Oxford: Oxford University Press. xv, 577 p. 2006.
- [Mil80] MILNE, James S.: * tale cohomology*. Princeton Mathematical Series. **33**. Princeton, New Jersey: Princeton University Press. XIII, 323 p. 1980.
- [Mil86] ———: *Abelian varieties*. Arithmetic geometry, Pap. Conf., Storrs/Conn. 1984, 103–150 (1986). 1986.
- [Mum70] MUMFORD, David: *Abelian varieties*. Oxford University Press. viii, 242 p. 1970.
- [RG71] RAYNAUD, Michel and GRUSON, Laurent: *Crit eres de platitude et de projectivit e. Techniques de « platification » d’un module*. In: *Invent. Math.*, **13** (1971), 1–89.
- [Sha86] SHATZ, Stephen S.: *Group schemes, formal groups, and  $p$ -divisible groups*. Arithmetic geometry, Pap. Conf., Storrs/Conn. 1984, 29–78 (1986). 1986.
- [Sza09] SZAMUELY, Tam as: *Galois groups and fundamental groups*. Cambridge Studies in Advanced Mathematics **117**. Cambridge: Cambridge University Press. ix, 270 p. 2009.
- [Tat66] TATE, John T.: *Endomorphisms of Abelian varieties over finite fields*. In: *Invent. Math.*, **2** (1966), 134–144.
- [Zar83] ZARKHIN, Yuri G.: *The Brauer group of an abelian variety over a finite field*. In: *Math. USSR, Izv.*, **20** (1983), 203–234.

TIMO KELLER, MATHEMATISCHES INSTITUT, UNIVERSIT AT BAYREUTH, 95440 BAYREUTH, GERMANY  
E-Mail address: *firstname.lastname@uni-bayreuth.de*